

MATHEMATICAL MODEL FOR THE 1.1 BILLION YEARS AGED SUN

Emil TATOMIR¹

Abstract

An algorithm for constructing evolutionary tracks for a star with the mass equal to one solar mass is given. The presented model can be applied to the stars belonging to the inferior main sequence, which have the proton-proton reaction as energy source and present a radiative core and a convective shell. This paper presents an original way of resolving the system of equations corresponding to the radiative nucleus by using Taylor's series in close vicinity to the center of the Sun. It also presents the numerical integration and the results for a 1.1 billion years aged solar model.

2010 *Mathematics Subject Classification*: 85-08 .

Key words: Mathematical model of sun, astrophysics

1 Basic formulae for the evolutive model

Consider that for the radiative core of the Sun the equations of hydrostatic equilibrium, mass distribution, luminosity and temperature are valid (see, e.g., Menzel and others, 1963; Aller and McLaughlin, 1965; Cox and Giuli, 1968), given respectively, given by:

$$\frac{dP(r)}{dr} = -\frac{GM(r)\rho(r)}{r^2}, \quad (1)$$

$$\frac{dM(r)}{dr} = 4\pi r^2 \rho(r), \quad (2)$$

$$\frac{dL(r)}{dr} = 4\pi^2 \rho(r) \varepsilon(r), \quad (3)$$

$$\frac{dT(r)}{dr} = -\frac{3}{4ac} \frac{\kappa(r)\rho(r)L(r)}{T^3(r)4\pi r^2}, \quad (4)$$

where $P(r)$, $M(r)$, $L(r)$ and $T(r)$ represent the values of the pressure, the mass, the luminosity and the temperature in a point placed at the distance r from the center of the star. By using Schwarzschild's (1958) transformations:

¹Faculty of Mathematics and Informatics, *Transilvania* University of Braşov, Romania, e-mail: emil.tatomir@gmail.com

$$\begin{aligned}
P(r) &= \frac{pGM^2}{4\pi R^4}, \\
M(r) &= qM, \\
L(r) &= fL, \\
T(r) &= t \frac{\mu H}{k} \frac{GM}{R}, \\
r &= xR,
\end{aligned} \tag{5}$$

the dimensionless variables p, q, f, t, x are introduced. With these variables, the system (1) – (4) becomes:

$$\begin{aligned}
\frac{dp}{dx} &= -\frac{pq}{tx^2}, \\
\frac{dq}{dx} &= \frac{px^2}{t}, \\
\frac{df}{dx} &= C \frac{px^2}{t}, \\
\frac{dt}{dx} &= -D \frac{pf}{t^4 x^2},
\end{aligned} \tag{6}$$

where we have denoted:

$$\begin{aligned}
C &= \frac{M}{L} (\varepsilon_{pp} + \varepsilon_{CN}), \\
D &= \frac{3Lk^4}{64\pi^2 ac \mu^4 H^4 G^4 M^4} \kappa, \\
A &= \frac{3Lk^4}{64\pi^2 ac H^4 G^4 M^4}.
\end{aligned} \tag{7}$$

The production of energy per gram-mass and per second due to the proton-proton reaction is given by the relation:

$$\begin{aligned}
\varepsilon_{pp} &= \varepsilon_0 \left(1 + 0.25 \rho^{\frac{1}{2}} T_6^{-\frac{3}{2}} \right) \left(1 + 0.012 T_6^{\frac{1}{3}} + 0.008 T_6^{\frac{2}{3}} + 0.00065 T_6 \right) \\
&\quad \rho X^2 10^6 T_6^{-\frac{2}{3}} e^{-33.804 T_6^{-\frac{1}{3}}},
\end{aligned} \tag{8}$$

where $\varepsilon_0 = 2.625$, ρ is the matter density expressed in $\frac{g}{cm^3}$, T_6 is the temperature expressed in $10^6 K$, while X denotes the hydrogen abundance.

Since the production of energy due to the carbon-nitrogen cycle will also be considered, this one is given by the relation:

$$\begin{aligned} \varepsilon_{CN} = & 7.94 \cdot 10^{27} \left(1 + 0.027T_6^{\frac{1}{3}} - 0.0037T_6^{\frac{2}{3}} - 0.00007T_6 \right) \\ & \left(1 + 1.75\rho^{\frac{1}{2}}T_6^{-\frac{3}{2}} \right) 0.00445\rho \cdot X^2 e^{-152.313T_6^{-\frac{1}{3}}} T_6^{-\frac{2}{3}}. \end{aligned} \quad (9)$$

The formulae (8) and (9) are in (Aller and McLaughlin, 1965). The total production of energy per gram-mass and per second is given by the formulae:

$$\varepsilon = \varepsilon_{pp} + \varepsilon_{CN}. \quad (10)$$

The opacity of stellar matter is considered to be given by the following analytical expressions:

- for $0.1 \leq T_6 < 10$:

$$\kappa = 0.19(1 + X) + a_1\rho(1 + X); \quad (11)$$

- for $10 \leq T_6 \leq 20$:

$$\kappa = 0.19(1 + X) + a_1\rho(1 + X) + \rho(1 - 0.1T_6)((1 + X)a_1 - b_1); \quad (12)$$

- for $T_6 > 20$:

$$\kappa = 0.19(1 + X) + b_1\rho; \quad (13)$$

where a_1 and b_1 have respectively the expressions:

$$\begin{aligned} a_1 = & 6.5 \cdot 10^4 \frac{Z(1 - 0.05T_6)}{T_6^2 + 2.5T_6^4} e^{-7.75\rho^{\frac{1+X}{T_6^3}}} + 4.15 \cdot 10^4 \left(\frac{X}{250T_6^4 - T_6^2} + \right. \\ & \left. \frac{1 - X - Z}{250T_6^4} \left(1 + 5.5e^{-|\frac{2}{3} - 2.873T_6|} \right) \right) e^{-2\rho^{\frac{1+X}{T_6^3}}} \end{aligned} \quad (14)$$

and

$$b_1 = \frac{35 + 75X + 190Z}{T_6^{3.5}} \quad (15)$$

in which Z features the metals.

For the convective shell of the Sun, the equations (1) and (2), and the adiabatic equation:

$$P(r) = K\rho^\gamma(r), \quad \gamma = \frac{5}{3}, \quad (16)$$

are considered to be valid. The ideal gas law is considered to be valid for the whole star.

Introducing the parameter:

$$(n+1)_{rad} = \frac{d \log P}{d \log T} = \frac{1}{D} \frac{qt}{pf}, \quad (17)$$

the radiative equilibrium is ensured by the condition:

$$(n+1)_{rad} > 2.5. \quad (18)$$

The system (1) – (4) is to be integrated with the boundary conditions (at center):

$$x = 0, \quad f = 0, \quad q = 0, \quad t = t_c, \quad p = p_c, \quad (19)$$

where t_c and p_c denote the dimensionless values of the temperature and pressure, respectively, at the Sun's center.

2 Numerical solution of the model

For the evolutive model which will be presented, we shall use the numerical results obtained by the author (Tatomir, 1986), which provide: pressure P , temperature T , dimensionless mass q and dimensionless luminosity f values in the points of a division:

$$\begin{aligned} x_1 < x_2 < \dots < x_{155}, \\ x_1 &= 0, \\ x_i &= (i-1)h, \end{aligned} \quad (20)$$

where the integration step was taken $h = 0.0058$.

The following values (corresponding to the parameters of the homogeneous model and the constants appearing in calculations) are used:

$$\begin{aligned} L_o^* &= 3.12E + 33 \left(\frac{erg}{s} \right), & c &= 2.9978E + 10 \left(\frac{cm}{s} \right), \\ R_o &= 6.96E + 10 (cm), & G &= 6.672E - 08 \left(\frac{cm^3}{gs^2} \right), \\ M_o &= 1.99E + 33 (g), & Q_{pp}^* &= 6.3E + 18 \left(\frac{erg}{s} \right), \\ k &= 1.379E - 16 (erg), & Q_{CN}^* &= 6.0E + 18 \left(\frac{erg}{s} \right), \\ H &= 1.672E - 24 (g), & X &= 0.709, \\ a &= 7.55E - 15 \left(\frac{dyne}{cm^2} \right), & Z &= 0.021, \end{aligned} \quad (21)$$

System (6) has a singularity in $x_1 = 0$; for calculating the values of p, q, f, t in the points x_1, x_2, \dots, x_7 , we have used in (14) the expanding in series method, obtaining:

$$p(x) = p_0 - \frac{1}{6} \frac{p_0^2}{t_0^2} x^2 + \left(\frac{1}{45} \frac{p_0^3}{t_0^4} - \frac{DC}{45} \frac{p_0^4}{t_0^8} \right) x^4 + 0 \cdot x^5 + \dots, \quad (22)$$

$$q(x) = \frac{1}{3} \frac{p_0}{t_0} x^3 + \left(\frac{DC}{30} \frac{p_0^3}{t_0^7} - \frac{1}{30} \frac{p_0^2}{t_0^3} \right) x^5 + 0 \cdot x^6 + \dots, \quad (23)$$

$$f(x) = \frac{C}{3} \frac{p_0}{t_0} x^3 + C \left(\frac{DC}{30} \frac{p_0^3}{t_0^7} - \frac{1}{30} \frac{p_0^2}{t_0^3} \right) x^5 + 0 \cdot x^6 + \dots, \quad (24)$$

$$t(x) = t_0 - \frac{DC}{6} \frac{p_0^2}{t_0^5} x^3 + \left(\frac{DC}{45} \frac{p_0^3}{t_0^7} - \frac{23D^2C^2}{360} \frac{p_0^4}{t_0^{11}} \right) x^4 + \dots, \quad (25)$$

where we have denoted $p_0 = p_c$ and $t_0 = t_c$.

By means of the values $X(x_i, 0)$, $\rho(x_i, 0)$, $T(x_i, 0)$ from the homogeneous model, one calculates the production of energy $\varepsilon_{pp}(x_i, 0) + \varepsilon_{CN}(x_i, 0)$ and the opacity $\kappa(x_i, 0)$, using the expressions (8) – (13).

As time step, we have chosen $\tau = 0.1 \cdot 10^9$ years; with this, the variation of chemical composition with time due to the nuclear reactions is given by:

$$X(x_i, \tau) = X(x_i, 0) - \left(\frac{\varepsilon_{pp}(x_i, 0)}{Q_{pp}^*} + \frac{\varepsilon_{CN}(x_i, 0)}{Q_{CN}^*} \right) \tau \quad (26)$$

At first for each integration point x_i and for the epoch τ the molecular weight μ and the values of the coefficients C, D are calculated:

$$\mu(x_i, \tau) = \frac{4}{3 + 5X(x_i, \tau) - Z}, \quad (27)$$

$$C(x_i, \tau) = \frac{1.99}{3.12} (\varepsilon_{pp}(x_i, 0) + \varepsilon_{CN}(x_i, 0)), \quad (28)$$

$$D(x_i, \tau) = A \frac{\kappa(x_i, 0)}{\mu^4(x_i, \tau)}, \quad (29)$$

where A is a numerical constant known from (7). In order to obtain the central values of density (ρ) and temperature (T) at instant τ , the nonlinear system:

$$\varepsilon_{pp}(0, 0) + \varepsilon_{CN}(0, 0) = \varepsilon_{pp}(\rho, T, X(0, \tau)) + \varepsilon_{CN}(\rho, T, X(0, \tau)), \quad (30)$$

$$\kappa(0, 0) = \kappa(\rho, T, X(0, \tau)), \quad (31)$$

is solved by means of the Newton-Kantorovici method.

Next we will show how the Newton-Kantorovici method is used for the evolutive solar model.

Introducing the next notations:

$$f(\rho, T) = \varepsilon_{pp}(\rho, T, X(0, \tau)) + \varepsilon_{CN}(\rho, T, X(0, \tau)) - \varepsilon_{pp}(0, 0) - \varepsilon_{CN}(0, 0), \quad (32)$$

$$g(\rho, T) = \kappa(\rho, T, X(0, \tau)) - \kappa(0, 0) \quad (33)$$

system (30) receives the form:

$$\begin{aligned} f(\rho, T) &= 0 \\ g(\rho, T) &= 0 \end{aligned} \quad (34)$$

And now

$$H : R^2 \rightarrow R^2, \quad H(\rho, T) = \begin{pmatrix} f(\rho, T) \\ g(\rho, T) \end{pmatrix} \quad (35)$$

where f and g are some functions of density ρ and temperature T .

Then system (33) can be written under the form:

$$H(\rho, T) = 0 \quad (36)$$

To obtain the solution of system (35) we start with an initial value of $\begin{pmatrix} \rho_k \\ T_k \end{pmatrix}$ which, in our case, is just the central value from the homogeneous model, $\begin{pmatrix} \rho_k \\ T_c \end{pmatrix}$. A better approximation of the solution of the system (33) can be obtained from the next formula:

$$\begin{pmatrix} \rho_{k+1} \\ T_{k+1} \end{pmatrix} = \begin{pmatrix} \rho_k \\ T_k \end{pmatrix} - \begin{pmatrix} \frac{\partial f(\rho_k, T_k)}{\partial \rho} & \frac{\partial f(\rho_k, T_k)}{\partial T} \\ \frac{\partial g(\rho_k, T_k)}{\partial \rho} & \frac{\partial g(\rho_k, T_k)}{\partial T} \end{pmatrix}^{-1} \cdot \begin{pmatrix} f(\rho_k, T_k) \\ g(\rho_k, T_k) \end{pmatrix} \quad (37)$$

We note with ρ_c^1 and T_c^1 the central values for the model of the type τ ; they are ρ^{k+1} and T^{k+1} in formula (36). Starting from ρ_c^1 and T_c^1 we obtain t_c^1 and p_c^1 from Schwarzschild's transformations (5) and from the law of gases

$$P(r) = \frac{1}{\mu} \frac{k}{H} \rho(r) T(r) \quad (38)$$

With $t_0 = t_c^1$ and $p_0 = p_c^1$ and by help of the series (22) – (25), we obtain the values for p, q, f, t in six points near the origin: x_2, x_3, \dots, x_7 . System (6) is integrated using the Adams-Bashforth method of the sixth order (Moszynski, 1973)

$$V_{k+1} = V_k + \frac{h}{1440} (4277f_k - 7923f_{k-1} + 9982f_{k-2} - 7298f_{k-3} + 2877f_{k-4} - 475f_{k-5}). \quad (39)$$

To improve the numerical results which have been obtained with the help of formula (38), the Adams-Moulton corrector method of the sixth order is used (Moszynski, 1973):

$$V_k = V_{k-1} + \frac{h}{1440} (475f_k + 1427f_{k-1} - 798f_{k-2} + 482f_{k-3} - 173f_{k-4} + 27f_{k-5}). \quad (40)$$

In this way we obtain the values for $t^1(x_i, \tau), p^1(x_i, \tau), f^1(x_i, \tau), q^1(x_i, \tau), \rho^1(x_i, \tau), T^1(x_i, \tau)$. The integration of the system(6) continues as long as $(n+1)_{rad} \leq 2.5$. The

integration of the model at the moment τ is repeated iteratively and we consider that for iteration n we have the next values:

$$X^n(x_i, \tau), \rho^n(x_i, \tau), T^n(x_i, \tau), p^n(x_i, \tau), f^n(x_i, \tau), q^n(x_i, \tau). \quad (41)$$

The passing from iteration n to iteration $n + 1$ of the model at moment τ is done in this way:

$$X^{n+1}(x_i, \tau) = X(x_i, 0) - \frac{1}{2}\tau \left(\frac{\varepsilon_{pp}(x_i, 0) + \varepsilon_{pp}^n(x_i, \tau)}{Q_{pp}^*} + \frac{\varepsilon_{CN}(x_i, 0) + \varepsilon_{CN}^n(x_i, \tau)}{Q_{CN}^*} \right) \quad (42)$$

$$\mu^{n+1}(x_i, \tau) = \frac{4}{3 + 5X^n(x_i, \tau) - Z} \quad (43)$$

$$C^{n+1}(x_i, \tau) = \frac{1.9891}{3.826} (\varepsilon_{pp}^n(x_i, \tau) + \varepsilon_{CN}^n(x_i, \tau)) \quad (44)$$

$$D^{n+1}(x_i, \tau) = A \frac{\kappa^n(x_i, \tau)}{(\mu^n(x_i, \tau))^4} \quad (45)$$

The central values of the model ρ_c^{n+1}, T_c^{n+1} at moment τ and at iteration $n + 1$ are obtained from the system:

$$\begin{aligned} \frac{1}{2} (\varepsilon_{pp}(0, 0) + \varepsilon_{pp}^n(0, \tau) + \varepsilon_{CN}(0, 0) + \varepsilon_{CN}^n(0, \tau)) = \\ \varepsilon_{pp}(\rho, T, X^{n+1}(0, \tau)) + \varepsilon_{CN}(\rho, T, X^{n+1}(0, \tau)) \end{aligned} \quad (46)$$

$$\frac{1}{2} (\kappa(0, 0) + \kappa^n(0, \tau)) = \kappa(\rho, T, X^{n+1}(0, \tau)). \quad (47)$$

The conditions to stop the iterations of the models at moment τ are:

$$|\rho_c^n - \rho_c^{n+1}| < \varepsilon_1 \quad (48)$$

$$|T_c^n - T_c^{n+1}| < \varepsilon_2 \quad (49)$$

When conditions (46) and (47) are accomplished it is considered that the model from iteration n is good and this model will be considered as being the one at moment τ .

The passing from a model at moment $m \cdot \tau$ to a model at moment $(m + 1) \cdot \tau$ is done in the identical way as the passing from the model at moment $\tau = 0$ to the model at moment τ .

3 Numerical results

Table 1 lists the numerical values featuring the solar model which corresponds to the age $\tau = 1.1 \cdot 10^9$ years.

Table 1

x	P	q	f	T	ρ	X
0.0	0.1885	0.0000	0.0000	14.2326	101.5883	0.5142
0.063	0.1650	0.0173	0.181	13.5863	90.3660	0.6765
0.116	0.1174	0.0905	0.6555	12.3873	71.2690	0.6919
0.162	0.773E-01	0.2063	0.9589	11.1588	51.7485	0.7015
0.208	0.467E-01	0.3499	1.1338	9.9796	34.9148	0.7065
0.307	0.143E-01	0.6486	1.1432	7.9395	13.0557	0.7088
0.406	0.408E-02	0.8716	1.1433	6.1412	4.5115	0.7089
0.510	0.977E-03	0.9830	1.1453	4.1726	1.4535	0.709
0.609	0.204E-03	0.9870	1.1463	2.4247	0.6725	0.709
0.707	0.302E-04	0.9889	1.1463	1.4550	0.1635	0.709
0.806	0.277E-05	0.9950	1.1463	0.9635	0.213E-01	0.709
0.893	0.383E-06	0.9989	1.1463	0.8943	0.3003E-02	0.709

The quantities appearing in Table 1 are:

- P - pressure (expressed in $10^{18} \frac{\text{dyne}}{\text{cm}^2}$);
- T - temperature (expressed in $10^6 K$);
- ρ - density (expressed in $\frac{g}{\text{cm}^3}$);
- X - hydrogen abundance;
- x - non-dimensional radius;
- q - non-dimensional mass;
- f - non-dimensional luminosity.

In order to compare the numerical data of the models obtained by the author to the observational data, we proposed plotting the model onto the $(\log \frac{L}{L_o}, \log T_{ef})$ - plane. In this respect we can represent our models on the observational diagram Hertzsprung-Russell. Since the energy is not produced in the region of the convective shell, namely $\varepsilon = 0$ and $L = \text{constant}$, the luminosity of the models is considered as given by:

$$L = L_o^* f(x_r) = L_o^* f(0.893) \quad (49)$$

The effective temperature is obtained from the well-known relationship valid for the black body:

$$L = 4\pi R^2 \sigma_R T_{ef}^4 \tag{50}$$

where $R = 6.96 \cdot 10^{10} cm$, while σ_R denotes the constant of Ştefan-Boltzmann: $\sigma_R = 5.6687 \cdot 10^{-5} \frac{erg}{cm^2 \cdot s \cdot deg}$.

Table 2 provides the values of the luminosity and effective temperature corresponding to the models obtained by the author.

Table 2

x	P	q	f	T	ρ
0.0	0.1885	0.0000	0.0000	14.2326	101.5883

In this table, the time τ is expressed in 10^9 years, f is (as previously) the non-dimensional luminosity, the luminosity L is expressed in $10^{33} \frac{erg}{s}$, and the effective temperature T_{ef} is expressed in K .

In figure 1 the model obtained by the author is plotted onto the Hertzsprung-Russell diagram. There are evolutionary models of stars of $1M_0$ belonging to Population I, corresponding to the epochs (ages) $\tau = 1.1 \cdot 10^9$ years. In comparison, the present-day Sun (aged about $4.5 \cdot 10^9$ years) is plotted on the diagram too, using the following data:

$$\begin{aligned} R_o &= 6.96 \cdot 10^{10} cm, \\ L_o &= 3.826 \cdot 10^{33} \frac{erg}{s}, \\ T_{ef_o} &= 5770K. \end{aligned} \tag{51}$$

Since there are not any other calculated models of the chemical composition on condition of $1.1 \cdot 10^9$ age and considered in this paper, not by any other author, the only thing that remains is to compare the model to the observational data.

The position of the $4.5 \cdot 10^9$ aged Sun on the $H - R$ diagram is presented in Figure 1, but the intermediate positions at different times can not be deduced in an observational way. The evolution of the Sun along the main sequence is not linear because the inner temperature increases in time, but the hydrogen abundance X decreases, modifying the values of the energy production and of the opacity given by the formulas (8) – (15). The position of the $1.1 \cdot 10^9$ aged model and of the actual Sun in the parallel positions with the main sequence and the distance between them, shows that the model which I have calculated approximates very well the position that the Sun would have occupied in the $H - R$ diagram when it would have had the age of $1.1 \cdot 10^9$ years.

The papers that had been quoted in the text were consulted in order to obtain this paper. The other papers quoted in the references are recommended to be read for a better understanding of the studied theme.




Figure 1: Position of the one billion years aged Sun

4 Personal contributions to the elaboration of an evolutive model

System (6), which has to be integrated on condition (19), presents an indetermination under the form of $\frac{0}{0}$. For the elimination of the indetermination we obtained the series (22) – (25).

We have shown how the Newton-Kantorovicz method can be used for the evolutive solar models and we have integrated system (6) using the method of successive approximations. The fact that the model, which has been calculated, gives good results from the comparison to the observation, means that this model placed in the $H - R$ diagram takes a correct position.

In conclusion, the original way of numerical solving, which is presented in section 3, can be used by all the evolutive models which have a radiative nucleus and a convective cover.

References

- [1] Aller, L.H., *Stellar structure*, University of Chicago Press, Chicago, 1965.
- [2] Cox, J.P., Giuli, R.T., *Principles of stellar structure*, Gordon and Breach, New York, 1968.

- [3] Christensen-Dalsgaard, J., Berthomieu, G., *Solar interior and atmosphere*, The University of Arizona Press, 1992.
- [4] Haselgrove, C.B., Hoyle, F., *A mathematical discussion of the problem of stellar evolution, with reference to the use of an automatic digital computer*. Mon. Not. Roy. Astr. Soc., **116** (1957), 515-526.
- [5] Kirsten, T. and the Gallex collaboration, *Inside the Sun*, Kluwer Academic Publishers, 1990.
- [6] Kurucz, R.L., *Stellar atmospheres, beyond classical models*, 1991.
- [7] Mendel, D.H., Bhatnagar, P.L., Sen, H.K. *Stellar Interiors*, Chapman and Hall, London, 1963.
- [8] Moszynski, K., *Numerical methods of solving the ordinary differential equations*, Technic Press, Bucharest, 1973.
- [9] Schwarzschild, M., 1958, *Structure and Evolution of the Stars*, Princeton University Press, Princeton, 1958.
- [10] Tatomir, E., *Thesis*, University of Cluj-Napoca, 1986.

